

Mouse Anti Clostridium Difficile Toxin B

#Cat: NB-47-04572-100UG Size: 0.1 mg

Description :	Mouse Anti Clostridium Difficile Toxin B
Specificity:	Clostridium Difficile Toxin B
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	5158
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.neo-biotech.com

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/20 - 1/200
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Bacterial
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide (NaN ₃)
Approx. Protein	IgG concentration 0.1mg/ml

Concentrations

External Database

Links

UniProt:

[P18177](#)

Synonyms

tcdB

Specificity

Mouse anti Clostridium difficile Toxin B antibody, clone 5158 recognizes *Clostridium difficile* toxin B. No reaction is observed with toxin A. *Clostridium difficile* is a gram-positive motile bacterium which is the leading cause of diarrhoea in developed countries. The incidence of disease in humans varies greatly with age, spore density and the administration of antibiotics.

Toxin B (TcdB) is a large exotoxin. Its role in disease is less well understood than that of toxin A, though it seems clear that TcdB contributes to disease. Both toxins modify the Ras superfamily of small GTPases via glycosylation, inactivating them and leading to the disruption of vital signaling pathways in the cell.

Further Reading

1. Voth, D.E. & Ballard, J.D. (2005) Clostridium difficile toxins: mechanism of action and role in disease. [Clin Microbiol Rev. 18 \(2\): 247-63.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040
available at: www.neo-biotech.com
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (NB-47-06061-500UG)

Rabbit Anti Mouse IgG (NB-47-05940-1ML)

Goat Anti Mouse IgG (NB-47-06055-500UG)

Goat Anti Mouse IgG IgA IgM (NB-47-06068-1MG)

Goat Anti Mouse IgG (NB-47-06060-1ML)

Goat Anti Mouse IgG (H/L) (NB-47-05906-500UG, NB-47-05908-500UG, NB-47-05909-500UG)

Rabbit Anti Mouse IgG (NB-47-05972-1MG)

Goat Anti Mouse IgG (Fc) (NB-47-05913-1MG, NB-47-05914-1MG)

Rabbit Anti Mouse IgG (NB-47-06079-1MG)